

A complete sample of radio sources in the North Ecliptic Cap, selected at 38 MHz – III. further imaging observations and the photometric properties of the sample

Mark Lacy^{1*}, Mary Elizabeth Kaiser^{2,3}, Gary J. Hill², Steve Rawlings¹
& Gareth Leyshon⁴

¹*Astrophysics, Department of Physics, Keble Road, Oxford, OX1 3RH.*

²*McDonald Observatory, University of Texas at Austin, TX78712-1083*

³*Bloomberg Center for Physics and Astronomy, Johns Hopkins University, 3400 Charles Street, Baltimore, MD21218*

⁴*Astronomy Dept, University of Wales College of Cardiff PO Box 913, Cardiff CF2 3YB*

1 February 2008

ABSTRACT

Further imaging observations of a sample of radio sources in the North Ecliptic Cap are presented and a number of new identifications are made. Using redshifts from spectroscopic data presented in a companion paper (Lacy et al. 1999b), the photometric properties of the galaxies in the sample are discussed. It is shown that: (1) out to at least $z \approx 0.6$ radio galaxies are good standard candles irrespective of radio luminosity; (2) for $0.6 \lesssim z \lesssim 1$ a large fraction of the sample have magnitudes and colours consistent with a non-evolving giant elliptical, and (3) at higher redshifts, where the R -band samples the rest-frame UV flux, most objects have less UV luminosity than expected if they form their stellar populations at a constant rate from a high redshift to $z \sim 1$ in unobscured star-forming regions (assuming an Einstein – de Sitter cosmology). The consequences of these observations are briefly discussed.

Key words: galaxies: active – radio continuum – galaxies: evolution – galaxies: formation

1 INTRODUCTION

Radio galaxies are at present the only high redshift galaxies which are not explicitly selected on the basis of high star formation rates, but which can nevertheless be found in reasonably large numbers. This makes them potentially very valuable for understanding the evolution of massive galaxies at early times. Unfortunately, the strong correlation of the radio and UV emission in the most radio-luminous galaxies (the “alignment effect”; see, e.g., Best et al. 1997 and Ridgway & Stockton 1997) indicates that their photometric properties are probably dominated by the presence of an AGN, at least at rest wavelengths $< 4000\text{\AA}$, a deduction which was confirmed by spectroscopically for most $z \sim 1$ 3C radio galaxies by Hammer, LeFèvre & Angonin (1993). This makes defining and studying samples of radio sources selected at fainter flux density levels important, as, due to the correlation of radio luminosity with the optical/UV luminosity of the AGN (e.g. Willott et al. 1999), such objects

should have only a very small fraction of their light produced by the AGN.

In two previous papers (Lacy, Rawlings & Warner 1992; Lacy et al. 1993; hereafter Papers I and II), we defined and presented radio and optical imaging data for a sample of radio sources selected from the 38-MHz survey of Rees (1990) in the region of the North Ecliptic Cap (NEC). In a companion paper (Lacy et al. 1999b; hereafter Paper IV) we present optical spectroscopy of this sample, along with that of a sample selected at 151MHz with a comparable flux limit and in the same region of sky (the 7C-III sample, Lacy et al. 1999c) and which therefore has most objects in common with this sample.

We have already discussed the outcome of high-resolution optical observations of a $0.5 < z < 0.82$ subsample of objects from the 7C-III sample which show that the radio – optical “alignment effect” in the 7C-III sample is weak, but probably present, even at wavelengths longward of $4000\text{\AA}$ (Lacy et al. 1999a). Thus is it possible that even in these radio-faint (≈ 20 -times fainter than 3C) samples, the radio source is influencing the properties of the host galaxy. As we discuss there, however, the scale of the alignments

* Guest observer, McDonald Observatory, University of Texas at Austin

does seem to change with radio luminosity, in the sense that the alignments of the 7C-III radio galaxies are detectable only on small scales (in ~ 15 kpc radius apertures), whereas the 3C radio galaxies have equally strong alignments at 15 kpc and 50 kpc. This could be because the highly radio-luminosity dependent mechanisms such as scattering of the hidden quasar nucleus and nebular emission disappear in the 7C-III objects, leaving a less luminosity-dependent mechanism which only operates on small scales. Relatively “passive” mechanisms may be responsible for these alignments, for example dust disks with axes along the radio axis and/or selection effects. If this were the case the photometric properties of the underlying galaxy would be unaffected by the presence of the radio source. However, it is possible that jet-induced star formation is only weakly radio-luminosity dependent, and if this mechanism operates then it would imply an intimate connection between the presence of the radio source and the stellar population of the host. The observational evidence for this mechanism is as yet only circumstantial however, and it is not clear whether such a mechanism could be effective (Icke 1999).

In this paper we discuss further optical, radio and near-infrared observations of the 8C sample, and use the spectroscopy presented in Paper IV to discuss the photometric properties and evolution of the radio source hosts. Unless otherwise stated we assume an $\Omega_M = 1, \Omega_\Lambda = 0$ (Einstein – de Sitter) cosmology with $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2 OBSERVATIONS

Radio snapshot observations of two of the sources in the sample, 8C1804+632 and 8C1826+660 were made with the VLA in C-array at 1.4 GHz on 1993 June 28 to attempt to resolve ambiguities of identification in Paper II. 8C1742+637 was observed in A-array at 1.4 GHz on 1995 June 29 for 2×2 min snapshots. These data were calibrated and analysed in the standard manner and the maps are presented in Fig. 1. In addition a four minute 8GHz A-array observation of 8C 1826+651 was made on 1996 December 31 to attempt to identify a central component. A 0.6 mJy point source was detected at 18 26 31.41 +65 10 46.0 (B1950), this is consistent with the identification in Paper II and corresponds to optical/infrared component ‘b’ of Fig. 2 being the probable radio source host (see also Lacy et al. 1998a).

Observations of several sources were made in the near-infrared *H*-band with ROKAM, a near-infrared camera with a 256×256 HgCdTe array, during 1993 May and June. ROKAM was placed at the f/18 straight Cass focus of the McDonald 2.7-m telescope, giving a pixel scale of 0.40 arcsec pixel $^{-1}$. Objects with estimated redshifts ≈ 1 in Paper II were the primary targets, though some lower redshift objects were also observed when the seeing was very poor. The observations are detailed in Table 1, and the photometry in Table 4. Greyscales of selected images are presented in Fig. 2. The observations were calibrated using the infrared standard stars HD136754 and HD162208 (Elias et al. 1982). Extinction was determined to be $0.15 \times$ airmass and a correction of 0.05 magnitudes was made for galactic extinction, assuming $E(B - V) = 0.085$ (Kolman et al. 1991) and the extinction curve of Cardelli, Clayton & Mathis (1992).

Further *R* and *I*-band observations of the unidentified

Table 1. *H*-band observations

Source	Date	Exposure time /s	airmass	Seeing /arcsec
8C1743+645	07/05/93	2970	1.22	2.8
	12/06/93	2970	1.22	1.9
8C1745+642	06/05/93	1890	1.20	1.6
	13/06/93	1620	1.20	1.5
8C1748+675	07/05/93	2160	1.32	2.4
8C1755+632	07/05/93	1890	2.0	2.7
8C1755+685B	07/05/93	1620	1.6	1.8
8C1757+653	09/05/93	1890	1.32	3.0
8C1803+661	08/05/93	1800	1.28	2.0
	12/06/93	5490	1.24	1.6
8C1805+635	11/06/93	3780	1.25	1.6
8C1807+685	07/05/93	2700	1.29	2.6
8C1811+633	08/05/93	2520	2.0	2.3
8C1815+682	09/05/93	1350	1.27	2.0
	12/06/93	3240	1.4	1.7
	13/06/93	810	1.36	1.5
8C1816+671	09/05/93	2610	1.28	2.3
8C1821+643	08/05/93	1080	1.4	3.2
8C1823+660	08/05/93	1980	1.6	3.5
8C1826+651	06/05/93	1620	1.22	1.6
8C1827+671	11/06/93	1080	1.4	1.5
	13/06/93	5670	1.25	1.4

sources in the sample were made on the McDonald 2.7-m using IGI (see Paper II) and on the William Herschel Telescope on La Palma using the Auxilliary Port. Details of these observations are given in Table 2. The extinction assumed at the WHT in July was $0.15 \times$ airmass in *R*, with a galactic extinction of 0.19 and the data were calibrated in the standard star PG 2331+055. In *I*-band, an extinction of $0.02 \times$ airmass was assumed for the observations made in 1994 January, and $0.05 \times$ airmass in 1994 July (higher due to Saharan dust). A value of $0.07 \times$ airmass was assumed for the McDonald observations. Galactic extinction in *I*-band was estimated to be 0.13 mag using the extinction curve of Cardelli et al. (1992).

3 ADDITIONS TO THE SAMPLE

Careful study of a 7C survey of the NEC (Lacy et al. 1995a) revealed four objects which, although absent from the original sample of Paper I due to confusion, should probably be included in the complete sample. Two of these remain excluded due to radio or optical confusion as discussed below. A full description of the objects including further radio maps

Table 2. Further optical observations

Source	Telescope/instrument	Filter	Date	Exposure time	airmass	seeing
				/s		/arcsec
8C 1743+645	McDonald 2.7-m/IGI	<i>R</i>	21/5/93	1200	1.20	2.0
8C 1748+670	WHT/Aux Port (TEK)	<i>R</i>	31/7/95	600	1.8	0.8
8C 1753+664	WHT/Aux Port (TEK)	<i>I</i>	25/1/95	900	1.7	0.8
8C 1754+643	WHT/Aux Port (TEK)	<i>I</i>	26/1/95	900	1.6	2.0
8C 1757+653	McDonald 2.7-m/IGI	<i>I</i>	21/5/93	3300	1.7	1.9
8C 1757+653	WHT/Aux Port (TEK)	<i>I</i>	25/1/95	600	1.6	0.8
8C 1814+670	WHT/Aux Port (TEK)	<i>I</i>	25/1/95	900	1.7	1.0
8C 1816+671	McDonald 2.7-m/IGI	<i>I</i>	21/5/93	2400	1.6	1.5
8C 1826+660A	WHT/Aux Port (TEK)	<i>R</i>	30/7/95	900	1.8	0.8

Table 3. Additions to and new identifications for objects in the 8C sample

Name	R.A. (1950)	Dec. (1950)	<i>R</i> -magnitude	Type	Notes
8C1732+672	-	-		?	Addition, but badly confused in radio so excluded from complete sample
8C1733+673	17 32 59.81	+67 19 32.3	22.6	G	Addition, <i>R</i> -magnitude from spectrum
8C1736+650	17 36 26.61	+65 04 10.7	23.0	G	Misidentification in Paper II, position uncertain, <i>R</i> from spectrum
8C1742+637	17 42 43.39	+63 46 11.8	22.6	G	Addition, identification assumed to be ‘a’ in Fig. 3
8C1743+645	17 43 28.55	+64 31 30.7	23.3	G	Misidentification in Paper II
8C1748+670	17 48 12.90	+67 03 54.2	23.9	G	Misidentification in Paper II ?
8C1754+643	17 54 11.89	+64 20 36.5	>23.1	G	Misidentification in Paper II. <i>I</i> = 21.1
8C1804+632	18 04 49.37	+63 13 07.9	~ 23	G	New identification; see Section 4
8C1805+635	18 05 37.33	+63 32 47.8	>23.7	G	Misidentification in Paper II
8C1814+670	18 14 16.09	+67 02 44.5	>22.9	G	New identification (<i>I</i> = 23.4)
8C1816+671	18 16 30.40	+67 11 00.2	>23.8	G	New identification (<i>I</i> = 22.4)
8C1826+660A	18 25 47.54	+66 02 24.0	23.8	G	New identification; note the finding chart in Paper II is incorrect
8C1827+652	18 27 59.13	+65 17 49.4	-	G?	Addition, but near bright stars so excluded from complete sample

Notes: This Table supplements table 2 of Paper II. Positions are the positions of the optical identifications (except for 8C1827+652 where it is the radio central component) and should be accurate to $\lesssim 1.5$ arcsec, except for 8C1736+650 whose identification is roughly at the mid-point of the radio hotspots, but is only seen in our spectrum and so the uncertainty is larger. Magnitudes are measured in the standard metric aperture of 63.9 kpc, if no redshift is known $z = 1$ is assumed.

and finding charts will be presented in Lacy et al. 1999c, but is summarised here and in Table 3 for completeness:

8C 1732+672 (7C 1742+6715)

The 1.5 GHz VLA map of this source is very confused, and so it has been temporarily removed from the sample again pending improved radio data.

8C 1733+673 (7C 1733+6719)

This is a small ($2.5''$) apparently double radio source, PA 15° . It is identified with an $R \approx 22.6$ galaxy (measured from the optical spectrum and corrected to the standard metric aperture).

8C 1742+637 (7C 1742+6346)

An FR II radio source with a marginal detection of a possible central component at 17 42 44.36 +63 46 15.0 (B1950) which is $51''$ in size (Fig. 1). The position of the nearest likely optical identification, ringed in Fig. 3, is given in Table 3. It lies along the radio axis, but not coincident with the putative central component, and has an *R*-magnitude of 22.6 in the standard metric aperture. Both the hotspots are projected close to what are probably foreground galaxies (the source lies 4.5 arcmin from the cluster Abell 2280) so significant gravitational lensing may be occurring.

8C 1827+652 (7C 1827+6517)

Another FR II radio source $17''$ in size at PA 155° . This object is very close to several bright stars so has again been temporarily removed from the sample.

4 COMMENTS ON INDIVIDUAL OBJECTS

8C1743+645

The identification given in Paper II (‘a’ in Fig. 2) closely coincides in position with the eastern radio hotspot. The *H*-band image (Fig. 2) reveals a faint red object coincident with the putative radio central component (Paper I) which is a much more likely identification. This galaxy is also just visible in *R*-band (Fig. 3). Lensing of the East hotspot by the (presumably) foreground galaxy ‘a’ is likely to be occurring.

8C1748+670

There are two candidate identifications, the first one was listed in Paper II, but our WHT *R*-band image (Fig. 3) also revealed a fainter object 1.4 arcsec closer to the radio central component whose position is given in Table 3. If real, this is the more likely identification. The revised position of the object discussed in Paper II is 17 48 12.95 +67 03 55.6 (B1950), the old position was in error, presumably due to the low signal:noise of the detection.




Figure 2. Selected H -band observations of the sources in the 8C sample. The host galaxies are indicated by either a ring around the radio position of Paper 1, or, where there are several candidate identifications, by lettering. Candidate companion objects are also indicated by lettering in some cases.

8C1754+643

The identification given in Paper II is incorrect; the true identification is a faint, diffuse galaxy located between the radio hotspots, ringed in Fig. 3.

8C1804+632

As discussed further in Paper IV, despite the additional radio imaging presented in Fig. 1, the nature of this radio source and its identification continues to elude us. The fairly compact source detected at 5GHz in Paper I now appears as just a part, perhaps a hotspot, of a larger, $\approx 30''$ radio source in our 20cm map. Spectra taken through objects 'a', 'F' and 'b' of Fig. 4 showed that all three were probably stars. The most likely identification at present is object 'c', which is unfortunately very close to star 'F'. Consequently the magnitude of this object is very uncertain, but we estimate $R \approx 23.6$ in a $2''$ diameter aperture.

8C1805+635

The original identification of this object in Paper II was an $R = 21.1$ galaxy 3 arcsec from line joining the midpoint of the radio hotspots ('a' in Fig. 2). The spectroscopy of Paper IV shows that the true identification is on the axis, and that 'a' is foreground, and possibly lensing the radio emission.

8C1814+670

There is a 3σ detection of a possible identification, 4.5 arcsec NW from the mid-point of the radio hotspots (Fig. 3).

8C1816+671

Another new identification, faintly visible in both H (Fig. 2) and R (Fig. 3).

8C1821+643

Better known as the quasar E1821+643, this object is one of the most luminous radio-quiet quasars (RQQ) known with $z < 0.5$, and thus has one of the brightest radio fluxes of an




Figure 3. New optical (*R*- or *I*-band) observations of sources in the 8C sample. The identification, or the best estimate of the radio source position where no identification exists, is indicated by the circle. The positions of the radio hotspots of 8C1742+637 are marked with stars.

RQQ. The optical, X-ray and near-infrared properties of the quasar have been studied by Kolman et al. (1993, 1991). The long-term variability study of Kolman et al. (1993) shows that the optical and UV flux from the quasar declined between 1987 and 1991; our *H*-band data is consistent with a decline from 1988 April 25, when Kolman et al. measured $H = 12.19$, to our observations in 1993 May 8 ($H = 12.6$),

although due to the errors in the photometry this result should be considered tentative.

4.1 8C1826+660

A further radio image of this object (Fig. 1) was taken to search for any association of the three radio sources dis-




Figure 1. New radio observations of sources in the 8C sample at 1.4 GHz: (a) 8C1804+632, (b) 8C1826+660 and (c) 8C1742+637. Contours are logarithmic, spaced by factors of two from ± 1 mJy/beam in (a) and (b), and by factors of $\sqrt{2}$ from 0.3 mJy/beam in (c). The restoring beam in (a) and (b) has FWHM $17'' \times 13''$ at PA 35° , that in (c) has FWHM $1.8'' \times 1.2''$ at PA 50° . The polarisation vectors in (c) are E -field and scaled such that 1-arcsec is equivalent to 1.1 mJy/beam. Negative contours are shown dashed.

cussed in Paper I. None was found, and henceforth they are treated as separate objects. A very faint identification has been found for 8C1826+660A (Fig. 3).

5 ANALYSIS

Table 4. H -band photometry of sources in the 8C sample


Source	H -magnitude	aperture diameter /arcsec	$R - H$
8C1743+645	18.8	5	4.7
8C1745+642	17.5	6	3.2
object ‘a’	18.9	5	> 4.2
object ‘b’	17.9	5	3.4
8C1748+675a	16.4	8.2	3.5
8C1748+675b	17.8	6	3.1
8C1748+675c	17.9	6	3.6
8C1755+632	15.3	18	2.6
8C1755+685B	16.9	6	3.2
8C1757+653	> 19.4	5	-
8C1803+661	18.3	7.1	2.8
8C1805+635	> 20.7	5	-
object ‘a’	18.5	4.6	2.6
8C1807+685	16.5	6	3.7
8C1811+633	16.2	7.1	3.1
8C1815+682a	17.8	6	3.3
8C1815+682b	19.6	4.6	1.7
8C1815+682c	17.1	6	3.8
8C1815+682d	17.0	6	3.0
8C1815+682e	17.9	6	3.8
8C1815+682f	17.7	6	3.7
8C1815+682g	18.3	4.6	3.4
8C1816+671	19.0	5	> 5.0
8C1821+643	12.6	8	-
8C1823+660	15.8	12	2.9
8C1826+651a	17.0	5	2.2
8C1826+651b	17.1	5	3.8
8C1826+651c	18.4	5	3.4
8C1826+651d	17.9	5	3.5
8C1827+671	18.5	8.2	2.6
object ‘a’	18.3	5	3.8
object ‘b’	17.1	8.2	3.1
object ‘c’	19.2	5	> 4.3
object ‘d’	18.9	5	> 4.6

5.1 The $R - z$ relation

In Fig. 5 we plot the $R - z$ relation for the 8C galaxies and quasars. Although such diagrams as this and the more famous $K - z$ relation are somewhat “blunt instruments” for studying radio galaxy evolution, where a number of different effects may be contributing to the light as a function of redshift, they have some useful features which can be exploited if they are used in conjunction with detailed studies of smaller samples of objects.

Table 5. Further optical photometry of sources in the 8C sample

Source	magnitude	aperture diameter /arcsec	$R - I$
8C1743+645	$R = 23.5$	5	-
8C1748+670	$R = 24.2$	3	-
8C1753+664	$I = 23.2 (3.0\sigma)$	5	> 0.4
8C1754+643	$I = 21.3$	5	> 1.3
8C1757+653	$I > 23.1$	5	-
8C1814+670	$I = 24.0 (3.0\sigma)$	2	> -0.8
8C1816+671	$I = 22.6 (3.7\sigma)$	5	> 2.6
8C1826+660A	$R = 24.0$	2	-



Figure 4. Finding chart for 8C1804+632. The objects mentioned in the text are labelled and the positions of the presumed radio hotspots from Fig. 1 marked with stars.

As expected, the scatter in the $R - z$ relation is fairly low (with a few notable exceptions) until $z \approx 0.6$, where the $4000\text{\AA}$ break moves through the R -band. Above this redshift there is a wide range in the amount of rest-frame UV light, as has been noted for the 3C sample (Lilly & Longair 1984), but generally the galaxies are less luminous than their 3C counterparts, consistent with the findings of Dunlop & Peacock (1993) for their PSR sample.

Up to at least $z = 1$ there seems to be an upper envelope to the magnitudes which traces the expected locus of an old stellar population, consistent with the idea that in many objects the host galaxy is not strongly evolving out to these redshifts. The presence of a weak alignment effect in the 7C sample (Lacy et al. 1998a) suggests some of the light may be influenced by the presence of the radio galaxy, however, so deep spectroscopy of these objects is really needed to confirm this.

Six objects have no redshift information, and are plotted as open squares at the top left of Fig. 5 with arbitrary redshifts. Their most likely redshifts are in the range $z = 1.2 - 1.8$ where no strong emission lines fall in the optical. They mostly have faint optical magnitudes and so do not affect the discussion below.

There is one interesting low- z outlier on this plot. 8C1743+637 ($R = 21.3$; $z = 0.324$) is in the cluster A2280,


Figure 6. $R - H$ versus redshift for a subset of the 8C sample. The objects plotted are those radio source identifications with $R - H$ -band colours in Table 4 and redshifts in Paper IV. Crosses represent narrow-line radio galaxies and filled circles broad-line radio galaxies. The dotted line represents the non-evolving 1Gyr starburst model of Fig. 5.

close to 8C1742+637, but was originally thought to be behind it from the point of view of both the radio structure (which is more FR II-like than FR I-like) and its magnitude (Paper II). It is possible that emission from a cluster galaxy is contaminating our spectrum, however.

5.2 $R - H$ colours

Fig. 6 shows the $R - H$ colours for the sample. The low redshift objects cluster around the no evolution prediction fairly well. There are two very red objects, 8C1743+645 with $R - H = 4.7$, but an unknown redshift, and 8C1816+671 with $R - H > 5$ which has a (tentative) redshift of 0.92 based on a single weak emission line (Paper IV). Presumably these are both $z \gtrsim 1$ objects with little blue light.

5.3 Discussion

A remarkable property of the radio galaxies out to at least $z \approx 0.6$ is their “standard candle” nature over a wide range in radio luminosities, illustrated here in Fig. 5 and also present in other low radio luminosity, moderate redshift samples (e.g. Rixon, Wall & Benn 1991). This has interesting implications for our understanding of what determines radio luminosity. To relate radio luminosity to the mass of the black hole, one could assume that radio jet power scales with accretion power and therefore with black hole mass in the Eddington limit; thus one might expect less radio-luminous galaxies to have less massive black holes. Correlations between black hole mass and the mass of spheroidal components in nearby galaxies (Kormendy & Richstone 1995, Magorrian et al. 1998), however, suggest that the small dispersion in host galaxy luminosities translates to a small dispersion in black hole mass. Two possible explanations which would allow similar black hole masses to produce a wide




Figure 5. The $R - z$ relation for the radio sources in the 8C sample, and the $z > 0.03$ objects from the mostly 3C low- z sample of Owen & Laing (1989). The magnitudes have been corrected to a standard metric aperture of 63.9 kpc according to the prescription of Eales et al. (1997). In the few cases where only an I -band magnitude has been measured (Table 5) an $R - I$ of 0.5, appropriate for a flat (in f_λ) spectrum has been assumed. The 8C objects are plotted as follows: solid squares represent narrow-line FRII radio galaxies, solid circles broad-line radio galaxies or quasars, and crosses represent FRI radio galaxies. The Owen & Laing sample has been plotted as triangles (FRII sources) or 3-pointed crosses (FRI sources). The cluster of open squares in the top left of the diagram represent 8C sources for which no redshift has been obtained, in these cases an aperture correction appropriate to $z = 1$ has been made. The vertical dashed line is at $z = 0.6$ where the $4000\text{\AA}$ break is in the middle of the R -band. The dotted line is the expected $R - z$ relation for a non-evolving galaxy formed in a single burst at $z = 20$ (using k -corrections from the Guiderdoni & Rocca-Volmerange galaxy models). The long-dashed line is the R -band magnitude expected from a star formation rate of $65 M_\odot \text{yr}^{-1}$ (see text). The solid line is the $R - z$ relation for 3C radio galaxies from Eales (1985).

range of radio luminosities are: (1) the radio jet power is constant for black holes of a given mass, and what determines the radio luminosity is the environmental density into which the jets propagate (Barthel & Arnaud 1996); or (2) the radio jet power is determined by a factor which operates within the “central engine” region of galaxies with only the most massive black holes and which has to be linked in some way to the accretion power to produce the radio-luminosity – emission-line luminosity correlation over a wide range in accretion rates (Willott et al. 1999). The lack of a strong correlation between radio luminosity and environment (Hill & Lilly 1991) and the correlation of radio-loud quasar optical and radio luminosities found by Serjeant et al. (1998) and

Willott et al. (1998) both argue that the dominant factor in determining the radio luminosity is jet power, although environmental effects may of course play a secondary rôle.

The excess blue light above that expected from a non-evolving elliptical seen in many objects at $z > 2$ could clearly have a number of origins, including ones associated with nuclear activity such as scattered quasar light and nebular continuum emission, but star formation is also a likely contributor. Indeed, if, as widely believed, radio galaxy hosts are giant ellipticals or their progenitors at all redshifts, then one can estimate the amount of UV light expected from star formation if radio galaxies form all their $\approx 3 \times 10^{11} M_\odot$ of stars in a single burst starting at $z = \infty$ and finishing at

$z = 1$. This is plotted on Fig. 5 as the long-dashed line, assuming a Scalo (1986) initial mass function. The average star formation rate, $65M_{\odot}\text{yr}^{-1}$ is high enough that *unobscured* star formation at or above this rate should be easily detected in our $z \sim 2$ objects. The fact that they all lie close to or fainter than this line could be attributed to several causes: (1) obscuration by dust may be extinguishing the UV light, (2) the star-forming episodes are of short duration, (3) the star-forming episodes occur mostly at very high redshift, (4) the star formation occurs in small sub-units before the radio galaxy host assembles, or (5) the cosmology is significantly different from Einstein – de Sitter.

Hughes and collaborators are performing a SCUBA-based submm survey of high redshift radio galaxies from the 3C and 6C (Eales 1985) surveys which should help to address point (1) (e.g. Hughes & Dunlop 1998; Hughes et al. 1998). Current limits on star formation rates are about an order of magnitude higher than $65M_{\odot}\text{yr}^{-1}$. Nevertheless, several detections of $2 \lesssim z \lesssim 4.3$ galaxies have already been made at this level. In addition Best et al. (1998) have reported a probable detection of the $z = 1.2$ radio galaxy 3C324. These detections, however, may only be upper limits too, in the sense that much of the FIR emission could be coming from dust heated by the AGN rather than from a starburst. The large dust masses involved, though, suggest at the very least that a large amount of star formation must have taken place in these objects by the time they are observed.

Possibility (2) can probably be ruled out unless the starbursts occur systematically before the formation of the radio source. Even so, such very bright starbursts should be easily detected in the IR/submm (if they are dusty) or in optical narrow-band searches if not.

Support for (3) or (4), very high redshift star formation either coherently or in small subunits which subsequently merge, comes from the studies of the old stellar populations in the radio galaxies 3C65 ($z = 1.2$; Lacy et al. 1995b; Stockton et al. 1995) and 53W091 ($z = 1.55$, Dunlop et al. 1996). The old stellar populations seen in some radio galaxies contrast with the studies of Zepf (1997) and Barger et al. (1999) who find relatively low numbers of very red galaxies in optical/infrared studies of the Hubble Deep Field (HDF) and its flanking fields. These show that only ~ 50 per cent of field ellipticals are in place by $z = 1$, and that the remainder probably formed at $z \lesssim 3$. The relatively high UV luminosity density produced by $z \sim 4$ Lyman-dropout galaxies found by Steidel et al. (1998), however, points to the possibility that at least some galaxies formed at higher redshifts. [Dust-obscured starbursts may affect this picture if they are important contributors to the star formation process in ellipticals, but currently this is uncertain (e.g. Trentham, Blain & Goldader 1998).]

A possible explanation for the apparent discrepancy between the ages of some radio galaxies and the field ellipticals is that giant elliptical galaxies formed earlier than normal $\sim L_*$ elliptical galaxies. This would be broadly consistent with an extrapolation of a trend deduced by Cowie et al. (1996) for more massive galaxies to have their major episodes of star formation at earlier epochs. If the oldest galaxies are the super- L_* population, they will be rare in the relatively small area field surveys made to date.

Explanation (5), an incorrect cosmology, remains possi-

ble. For example an $\Omega_M = 0.5$, $\Omega_\Lambda = 0.5$ cosmology with the same value of H_0 only requires a minimum star formation rate of $\approx 50M_{\odot}\text{yr}^{-1}$ to form all the stars by $z = 1$, and the emission will also appear fainter at a given redshift due to the higher luminosity distance. These combine to make the R -magnitude for a $z = 2$ galaxy forming stars at the minimum rate 0.8 mag fainter, for example.

The lack of UV emission also argues somewhat against jet-induced star formation being an important contributor to the star formation process of the galaxy as a whole, at least at $z \lesssim 3$. If it were, one would expect all radio galaxies to show large amounts of UV and/or submm emission. It remains possible though that an initial episode of star formation activity could be triggered by an early outburst of radio jet activity at high redshift, and that subsequent episodes of radio jet activity would produce more modest starbursts, consistent with the relatively low limits on the unobscured star formation rates we infer for the 8C radio galaxies.

ACKNOWLEDGEMENTS

We thank the staff at the McDonald Observatory and the WHT for their assistance. The WHT is operated on the island of La Palma by the Royal Greenwich Observatory in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. The VLA is operated by Associated Universities Inc. under a cooperative agreement with the National Science Foundation.

REFERENCES

- Barger A.J., Cowie L.L., Trentham N., Fulton E., Hu E.M., Songaila A., Hall D., 1999, AJ, 117, 102
- Barthel P.D., Arnaud K.A., 1996, MNRAS, 283, 45
- Best P.N., Longair M.S., Röttgering H.J.A., 1997, MNRAS, 292, 758
- Best P.N., Röttgering H.J.A., Bremer M.N., Cimatti A., Mack K.-H., Miley G.K., Pentericci L., Tilanus R.P.J., van der Werf P.P., 1998, MNRAS, in press (astro-ph/9809284)
- Cardelli J.A., Clayton G.C., Mathis J.S., 1989, ApJ, 345, 245
- Connolly, A.J., Szalay, A.S., Dickinson, M., Subbarao, M.U., Brunner, R.J. 1997, ApJ, 486, 11.
- Cowie L.L., Songaila A., Hu E.M., Cohen J.G., 1996, AJ, 112, 839
- Dey, A., van Breugel, W., Vacca, W., Antonucci, R. 1997, ApJ, 490, 698
- Dunlop, J.S., & Peacock, J.A. 1993, MNRAS, 263, 936
- Dunlop, J.S., Peacock, J.A., Spinrad, H., Dey, A., Jiminez, R., Stern, D. & Windhorst, R. 1996, Nat., 381, 581
- Elias J.H., Frogel J.A., Matthews K., Neugebauer G., 1982, AJ, 87, 1029
- Eales S.A., 1985, MNRAS, 217, 179
- Eales S.A., Rawlings S., Law-Green D., Cotter G., Lacy M., 1997, MNRAS, 291, 593
- Hammer, F., Le Fevre, O., & Angonin, M.C. 1993, 362, 324.
- Hill G.J., Lilly S.J., 1991, ApJ, 367, 1
- Hughes D.H., Dunlop J.S., to appear in Carilli et al., eds, "Highly Redshifted Radio Lines", NRAO, Greenbank 9-11 October 1997. PASP, (astro-ph/9802260)
- Hughes D.H., Dunlop J.S., Archibald E.N., Rawlings S., Eales S.A., 1998, to be published in the proceedings of "The Birth of Galaxies", Xth Rencontres de Blois (astro-ph/9810253)

- Icke V., 1999, in, Röttgering H.J.A., Best P.N., Lehnert M.D., eds, "The most distant radio galaxies". Royal Netherlands Academy of Arts and Sciences, Amsterdam, p. 217
- Kolman M., Halpern J.P., Shrader C.R., Filippenko A.V., 1991, ApJ, 373, 57
- Kolman M., Halpern J.P., Shrader C.R., Filippenko A.V., Fink H.H., Schaeidt S.G., 1993, ApJ, 402, 514
- Kormendy J., Richstone D., 1995, ARA&A 33, 581
- Lacy M., Rawlings S., Warner P.J., 1992, MNRAS, 256, 404 (Paper I)
- Lacy M., Hill G.J., Kaiser M.E., Rawlings S., 1993, MNRAS, 263, 707 (Paper II)
- Lacy M., Rawlings S., Eales S., Dunlop J.S., 1995b, MNRAS, 273, 821
- Lacy M., Riley J.M., Waldrum E.M., McMahon R.G., Warner P.J., 1995a, MNRAS, 276, 614
- Lacy M., Ridgway S.E., Wold M., Lilje P.B., Rawlings S., 1998a, MNRAS, in press
- Lacy M., Rawlings S., Hill G.J., Bunker A.J., Stern D., Ridgway S.E., 1998b, in press (Paper IV)
- Lacy M., Blundell K.M., Hill G.J., Kaiser M.E., Rawlings S., 1999c, in preparation
- Lilly, S.J. & Longair, M.S. 1984, MNRAS, 211, 833.
- Madau, P. et al. 1996, MNRAS, 283, 1388
- Magorrian J., Tremaine S., Richstone D., Bender R., Bower G., Dressler A., Faber S.M., Gebhardt K., Green R., Grillmair C., Kormendy J., Lauer T., 1998, AJ, 115, 2285
- Owen F.N., Laing R.A., 1989, MNRAS, 238, 357
- Rees N., 1990, MNRAS, 244, 233
- Ridgway S.E., Stockton A.N., 1997, AJ, 114, 511
- Rixon G.T., Wall J.V., Benn C.R., 1991, MNRAS, 251, 243
- Scalo J.M., 1986, Fund. Cosmic Phys., 11, 1
- Serjeant, S., Rawlings, S., Maddox, S.J., Baker, J.C., Clements, D., Lacy, M. & Lilje, P.B. 1998, MNRAS, 294, 494
- Steidel C.C., Adelberger K.L., Giavalisco M., Dickinson M., Pettini M., 1998, ApJ, in press (astro-ph/9811399)
- Stockton A.N., Kellogg M., Ridgway S.E., 1995, ApJ, 443, L69
- Trentham N., Blain A.W., Goldader J., 1998, MNRAS, in press
- Willott C.J., Rawlings S., Blundell K.M., Lacy M., 1998, MNRAS, 300, 625
- Willott C.J., Rawlings S., Blundell K.M., Lacy M., 1999, MNRAS, in press
- Zepf S.E., 1997, Nat, 390, 377